




Indicator 1.1: Percentage of children aged 6 - 59 months, with Stunting (a low height for age < -2 standard deviations (SD) of the WHO Child Growth Standards median)...




Indicator.1.1.1: Stunting by Wealth Quintile



Indicator. 1.1.2: Stunting by Age




Indicator 1.1.3: Stunting by Sex




National Nutrition Profile


Indicator 1.3: Percentage of children under five years with a low weight for age (< -2 standard deviations (SD) of the WHO Child Growth Standards median)




Indicator 1.2.1: Wasting by Wealth Quintile




Indicator 1.2.2: Wasting by Age




Indicator 1.2.3: Wasting by Sex




Indicator 1.3: Percentage of children under five years with a low weight for age (< -2 standard deviations (SD) of the WHO Child Growth Standards median)




Indicator 1.3.1: Underweight by Wealth Quintile







Indicator 1.3.2: Underweight by Age




Indicator 1.3.3: Underweight by Sex







Indicator 1.8: Percentage of children aged 0-23 months who were put to the breast within one hour of birth




Indicator 2.1: percentage of children under one year with low birth weight (a weight at birth <2,500grams)




Indicator 2.2: Percentage of Underweight Women, 15 to 49 years of age (with a Body Mass Index (BMI) less than 18.5)




Indicator 2.3: Percentage of Underweight Adolescents, 10 – 19 years of age (with a Body Mass Index (BMI) less than 18.5)




Indicator 2.4: Percentage of Overweight Women, 15 – 49 years of age (with a Body Mass Index (BMI) more than 24.9)



Indicator 2.5: Percentage of overweight adolescents of 10-19 years



Indicator 3.1: Prevalence of Anaemia among Children aged 6 – 59 Months (with a haemoglobin concentration < 110g/l at sea level)




Indicator 3.1.1: Anaemia by Wealth Quintile




National Nutrition Profile


Indicator 3.1.2: Anaemia by Age



Indicator 3.1.3: Anaemia by Sex



Indicator 3.2: Prevalence of Anaemia in pregnant women (with a haemoglobin concentration <110g/l at sea level)



Indicator 3.3: Percentage of children aged 6-59 months who received one or two doses of vitamin A supplements within one year

